

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS


IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**


1/61

Figure 1

Fusion of Cochleate Membrane with Target Cells



Phase/Fluorescent Image




Fluorescent Image

2/61

Figure 2

Formulation of Hydrophobic Cargo Moiety Into Cochleates: Solvent Drip Method




3/61

Figure 3


Formulation of Hydrosoluble Cargo Moieties Into Cochleates:

Trapping method with preformed liposome




4/61

Figure 4




5/61

Figure 5




6/61

Figure 6




7/61

Figure 7




8/61

Figure 8




9/61




10/61

Figure 10
Survival Data




11/61

Figure 11
Comparison of Bacterial Burden for *in vivo* Study of AmB formulations




12/61




13/61


Figure 13



Re-suspended AmB Cochleates with 1.28% V-E (AmB to V-E w/w)




Re-suspended AmB Cochleates with 1.28% V-E (AmB to V-E w/w) and After Adding EDTA



14/61

Figure 14


SURVIVAL DATA



15/61

Figure 15


Comparison of Bacterial Burden for in vivo Study of AmB formulations



16/61

Figure 16


In Vitro Efficacy of AmB Cochleates



17/61

Figure 17


Survival Data



18/61

Figure 18


Efficacy of CAMB formulations VS AmB/deoxycholate



19/61


Figure 19

Kinetics of Tyrophostin in Cochleates vs. Free




20/61

Figure 20



21/61


Figure 21



22/61


Figure 22

ZnTPP In solution in 100% DMSO and in Cochleates



23/61


Figure 23



24/61

Figure 24

Cochleates using regular trapping method




25/61


Figure 25

ZnTPP in cochleates

After one hour



After 24 hours




26/61


Figure 26

ZnTPP in solution in DMSO

After one hour



After 24 hours




27/61


Figure 27

Cochleates containing Pyrene DOPE

After 1 hour



After 24 hours




28/61


Figure 28

Cochleates containing Pyrene DOPE and ZnTPP

After 1 hour



After 24 hours



29/61




Figure 30






Figure 31



30/61

Figure 32



31/61

Fluorescence



Figure 33A

In vitro uptake of Rho-PE-lipid
precipitates

Lipid Precipitates Without
Aggregation Inhibitor

Phase Contrast and Fluorescence





Figure 33B

In vitro uptake of Rho-PE-lipid
precipitates

Lipid Precipitates With
Aggregation Inhibitor



32/61

Figure 34B

Lipid Precipitates Without Aggregation Inhibitor





Figure 34A


Lipid Precipitates With Aggregation Inhibitor



33/61

Figure 35

A. Rhodamine Cochleates With Half and Half



B. Rhodamine Cochleates With Whole Milk



C. Rhodamine-Cochleates With Evaporated Fat Free Milk




D. Rhodamine Cochleates Without Aggregation Inhibitor




34/61

Figure 36

A: Rhodamine Cochleates Prior to
Addition of Milk




B: Rhodamine Cochleates With Milk



35/61

Figure 37A




Acetaminophen Lipid Precipitates With Casein



Acetaminophen Lipid Precipitates Without Casein

Figure 37B



Aspirin Lipid Precipitates With Casein




Aspirin Lipid Precipitates Without Casein

36/61

Figure 38


Comparison of *In Vivo* Efficacy of Aspirin Formulations in Mice (Right Hind Paw)



37/61


Figure 39

Gastric Irritation in Rat Model




38/61

Figure 40



39/61

Figure 41
In Vitro Efficacy of Anti-Inflammatory Cochleates



40/61

Figure 43






Figure 42



41/61


Figure 44




42/61

Figure 45
Vancomycin


A. Liposomes




B. Precipitates with casein



D. Precipitates w/o Casein
upon addition of EDTA




C. Precipitates without Casein



43/61

Figure 46


Vancomycin Precipitate Efficacy @ 3hrs



In Vitro Efficacy of Vancomycin Precipitates in Macrophages Infected with *Staphylococcal aureus* at 3 hours post infection.

Figure 47

Vancomycin Precipitates Efficacy @ 6hrs




In Vitro Efficacy of Vancomycin Precipitates in Macrophages Infected with *Staphylococcal aureus* at 6 hours post infection.

45/61


Figure 48

Tobramycin


A. Liposomes




C. Precipitates without Casein



B. Precipitates with casein




D. Precipitates w/o Casein upon addition of EDIA



46/61

Figure 49


Efficacy of Tobramycin Formulations Against
Pseudomonas aeruginosa at 3 Hours Post-Infection



47/61


Figure 50

Efficacy of Tobramycin Formulations Against
Pseudomonas aeruginosa at 6 Hours Post-Infection




48/61

Figure 51




49/61

Figure 52



50/61


Figure 53



51/61

Figure 54

Particle size distribution of caspofungin lipid precipitates before
and after the addition of BSA



52/61

Figure 55






Figure 56



54/61

Figure 57


Structure of caspofungin lipid precipitates as a function of pH.




pH=1




pH=6




pH=8



pH=4



pH=7









pH=9

55/61

Figure 58

AmB Cochleates Containing 0.2% (w/w) Parabens




0.1% (w/w) Methylcellulose

0.3% (w/w) Methylcellulose


0.5% (w/w) Methylcellulose

Figure 59



57/61

Figure 60



58/61

Figure 61

In vitro Efficacy of J774A






Figure 62



60/61

Figure 63



61/61

Figure 64

